Perhaps I know best why it is man alone who laughs; he alone suffers so deeply that he had to invent laughter.
-Nietzsche, The Will to Power

Introduction

People often indulge in dark humor, a psychological state evoked by jokes with immoral or repulsive content (Bloom, 2010). Why do people enjoy making light of grotesque topics such as death or sex? In this paper, I argue that while dark humor violates moral principles, the dark humorous state is a necessary source of chemical pleasure.

Background

1.1. Humor

One usually enters a humorous state through incongruency detection and resolution. In other words, individuals recognize an event's inconsistency with their expectations, thus evoking surprise and a piecing-together process, all to find a joke funny (Suls, 1972). For example, why can't a bicycle stay standing? It feels "tired." To break down this joke, "tired" is a play on two definitions: a bicycle's tires and the human feeling of being tired. The "aha!" moment is the resolution that one arrives at after reconstructing this joke in their head.

Be that as it may, individuals more frequently indulge in atypical humor, which is evoked by ideas that cannot be pieced together (Ruch, 1992). For instance, people may find it hilarious to witness a dog using curse words towards its owner. Realistically, dogs cannot speak like humans do, let alone use vulgar language. This incongruency can only be resolved by recognizing absurdity of the situation, rather than reconstructing the joke through logical means.

Dark humor mixes immorality into the typical and atypical. Sexual jokes commonly evoke the typical dark, humorous state since the mention of sex outside of intimate settings is incongruent with morals (Russell, 1999). To demonstrate, what is the first thing a man puts in a woman when they get married? The wedding ring! The incongruency of this joke lies between the question's suggestive nature and the actual answer. The question's phrasing intends to motivate an answer related to sexual intercourse, but the answer defies that expectation by providing a literal rather than suggestive answer, the wedding ring. On the dark side of atypical humor, people joke about ideas that evoke repulsive feelings in most people, such as having sexual motivations toward their dog. This example, like many other dark jokes, transcends the absurdity of the normal atypical humorous state by violating moral principles.

It is quite natural to question why people indulge in these psychological states. However, one must comprehend the neurobiological grounds for humor and its behavioral expression before understanding the benefits of jesting at such moral defance.

1.2. Chemical Motivation

Humor and its ensuing laughter induce the release of a brain chemical called dopamine, which exerts pleasurable sensations. More specifcally, humorous laughter activates the mesolimbic pathway, a dopamine-abundant brain circuit that essentially motivates behaviors for obtaining sources of pleasure (Ikemoto, 2010). Azim and colleagues conducted fMRI studies on a sample of 10 men and 10 women. During the scans, each participant was presented with a cartoon, and they pressed a button if they found it funny. Subsequently, the subjects ranked each cartoon's hilarity on a 1-10 scale. The researchers compared fMRI results with participants' comedic ratings to determine whether blood fow changes in specifc brain structures corresponded with funniness. Consequently, participants exhibited higher blood fow in the nucleus accumbens, a key component of the mesomvm[key co6 of the000005200rw052005ay) 0 e0rw05_,

higher amount of stimulation, the patient's display of positive emotion increased as a smile grew into laughter. MRI and CT scans showed activation patterns in a set of connections called the medial forebrain bundle (MFB), which regulates behaviors that approach sources of pleasure (Coenen et al., 2022). One of the MFB's offshoots, the superolateral branch, converges with key structures of the mesolimbic pathway (Coenen et al., 2009). These results show that laughter coincides with the activation of the MFB, which includes connections with the dopaminergic pleasure pathway. Together with the fndings of the previous study, dopaminergic release within the mesolimbic pathway likely rewards humor evocation and induces the humorous execution of laughter.

1.3. Chemical Delight and Relaxation

Serotonin

Moreover, humorous laughter elevates levels of serotonin, the brain molecule implicated in mood regulation (Mohammad-Zadeh et al., 2008).

The effects of humor on serotonergic release are best perceptible through changes in depressive symptom severity. 17 elderly partici-

pamine that combats pain with pleasure (Miller et al., 2014). Research supports the notion that the effects of beta-endorphin result from humor induction. 68 patients receiving hemodialysis, a procedure that filters the blood to compensate for kidney failure (Elliott, 2000), participated in sixteen 30-minute biweekly laughter yoga sessions. During this exercise, participants were instructed to tell jokes to each other while forcing their laughter so they could eventually produce natural laughter. The researchers collected blood samples to measure beta-endorphin levels before the first session and after the last session. As a result, patients had lower blood endorphin levels after all 16 sessions (Özer & Ateş, 2021). Based on these findings, humorous laughter, even when forced, increases the release of beta-endorphin. This increase in concentration causes heightened dopaminergic levels that evoke a pleasant counter to pain signals.

Cortisol

Additionally, humor-induced laughter reduces levels of cortisol, a hormone that circulates throughout the body to modulate chronic stress in various contexts, including endurance activities, shift work, major life events, and emotional hardship (Staufenbiel et al., 2013). One study examined the relationship between cortisol and humorous laughter. 120 healthy university students participated in one of three activities: laughter yoga, comedy movie, and reading. Researchers collected salivary samples immediately before, immediately after, and 30 minutes after participants performed their assigned activity. In effect, the students who engaged in laughter yoga and the comedy movie consisted of lower salivary cortisol levels than those who read a book (Fujisawa et al., 2018). As implied by the results, laughter alleviates stress through the reduction of cortisol levels**.**

1.4. Chemical Sociality

Another brain molecule, oxytocin, motivates group laughter. The brain circuit associated with social behavior consists of numerous nodes that are abundant in neuropeptides (proteins associated with brain function) (Albers, 2015). One of these proteins, called oxytocin (OT), influences prosocial behaviors (behavior that benefits others), including interpersonal attachment (Jurek & Neumann, 2018). Recent studies demonstrate that the induction of humor and laughter are oxytocin-mediated and, therefore, prosocial behaviors. Participants viewed a brief clip of a comedy television show with a close friend after receiving intranasal OT or a placebo (a substance that has no effect). Consequently, the group that received OT during the show laughed more than those who consumed the placebo. This evidence substantiates that laughter is a prosocial behavior and that oxytocin promotes social laughter. Unlike other brain chemicals involved in humor, oxytocin concentrations did not change in response to laughter; rather, they induced the humorous state to eventually produce laughter. Therefore, oxytocin likely contributes to the initiation rather than the reinforcement of group laughter.

Considering dopamine's involvement in pleasure motivation through the mesolimbic pathway (Ikemoto, 2010), oxytocin likely interacts with dopamine-abundant brain regions to prompt social laughter. Hung and colleagues performed multiple experiments on rat models to test this notion. In one of the procedures, the control group remained untouched while the other received an injection to inhibit neurons within an oxytocin-abundant structure called the paraventricular nucleus (PVN). These neurons communicate with the ventral tegmental area (VTA), a key structure of the mesolimbic pathway. 2 weeks after inhibition; the researchers performed a socially conditioned place preference (CPP) test to determine whether the rats prefer to stay in an environment associated with social interaction. As a result, rats that received inhibitory injections to the PVN spent significantly less time in the social context than the control group. These experimental findings suggest that oxytocinergic connections must interact with pleasure motivation circuitry to elicit social behavior. Since the former study conveys humorous laughter as a social behavior, collective humorous expression likely follows this same neurobiological mechanism.

Furthermore, endorphinergic release reinforces oxytocin-dopamine-mediated laughter within a social setting. One 2017 study confirms this notion. Twelve healthy male participants engaged in a social laughter manipulation experiment. Each subject spent 30 minutes in a room without any social contact. Then, they provided baseline brain activity via PET scans. Subsequently, participants spent another 30 minutes watching a pre-arranged series of comedic YouTube clips with two close friends. A video camera recorded every session while a microphone tracked the number of laughter outbursts. The researchers conducted another set of PET scans immediately after social laughter, and they quantified the number of micro-opioid receptors (MORs) (including endorphinergic receptors) in various brain regions. Relative to baseline, PET images after viewing the clips with close friends showed a higher release of endorphins and an increased presence of MORs in structures that comprise the dopaminergic pleasure pathway. As supported by the results, laughter increases the availability of MORs to allow more endorphins to bind and induce dopaminergic release. This gives rise to pleasurable sensations that counteract pain and reinforce laughter during social interaction.

In the rest of this paper, I utilize the evidence of laughter's chemical pleasures, as revealed by neuroscience, to explain why dark humor is a necessary source of reward. I initially argue that dark humor is a gratifying experience, and then I assert that this psychological state combats distressing feelings such as pain or stress. Subsequently, I refute possible counterarguments to further support my assertion. Finally, I provide caveats and concluding remarks.

Dark Humor is the Brain's Necessary Evil

Despite its morally defiant nature, dark humor allows the brain to release chemicals that create pleasurable sensations. One study provides evidence for this notion. Thirty male volunteers viewed 32 single-frame sexual humor cartoons while under an fMRI. The procedure consisted of two phases: anticipation and outcome. During the anticipatory phase, the cartoon appeared for 1000 ms and ensued with a 2500 ms anticipatory delay period. Subsequently, the outcome phase required participants to determine whether a target number was smaller or larger than 5. They witnessed the correctness of their answer and then received either an erotic, monetary, or non-reward outcome. Finally, participants gave a rating pertaining to their en3758 Tm[pertaining)-0.8 (to zo their ertaining3763.6(,)0.5 8